

Conformal self-dual fields

R.R. Metsaev*

*Department of Theoretical Physics, P.N. Lebedev Physical Institute,
Leninsky prospect 53, Moscow 119991, Russia*

Abstract

Conformal self-dual fields in flat space-time of even dimension greater than or equal to four are studied. Ordinary-derivative formulation of such fields is developed. Gauge invariant Lagrangian with conventional kinetic terms and corresponding gauge transformations are obtained. Gauge symmetries are realized by involving the Stueckelberg fields. Realization of global conformal symmetries is obtained. Light-cone gauge Lagrangian is found. Also, we demonstrate use of the light-cone gauge for counting of on-shell degrees of freedom of the conformal self-dual fields.

* E-mail: metsaev@lpi.ru

1 Introduction

In Poincaré and conformal supergravity theories, the self-duality manifests itself in different ways. In Poincaré supergravity theories, some of the antisymmetric tensor fields are not self-dual, while their field strengths are self-dual (see, e.g., Ref.[1]).¹ In contrast to this, in conformal supergravity theories, some of the antisymmetric tensor fields are self-dual themselves, while their field strengths are not self-dual (see, e.g., Ref.[6]).

In Poincaré supergravity theories, the antisymmetric tensor fields are realized as gauge fields, while in the standard approach to conformal supergravity theories there are no gauge symmetries related to the self-dual antisymmetric tensor fields. Note also that the antisymmetric tensor fields of Poincaré supergravity theories describe ghost-free dynamics, while the ones of conformal supergravity theories contain of ghost degrees of freedom.

In this paper we discuss the self-dual antisymmetric tensor fields of conformal supergravity theories (which are well defined only in $d = 4, 6$) and their counterparts in space-times of arbitrary even dimensions. It is these self-dual antisymmetric tensor fields that will be referred to as conformal self-dual fields, or shortly as self-dual fields in this paper.

The standard formulation of the self-dual fields involves exotic kinetic terms². These exotic kinetic terms can be reexpressed in terms of kinetic terms involving the standard D'Alembertian operator but this leads to higher derivatives (see e.g. Ref.[6]). Also, as was mentioned above, in the standard approach, there are no gauge symmetries associated to such self-dual fields.

The purpose of this paper is to develop an ordinary (not higher-) derivative, gauge invariant, and Lagrangian formulation for the self-dual fields³. In this paper, we discuss free self-dual fields in space-time of even dimension $d \geq 4$. Our approach to the self-dual fields can be summarized as follows.

i) We introduce additional field degrees of freedom (D.o.F), i.e., we extend space of fields entering the standard formulation of self-dual fields. These additional field D.o.F are supplemented by appropriate gauge symmetries⁴. We note that these additional field D.o.F are similar to the ones used in the gauge invariant formulation of massive fields. Sometimes, such additional field D.o.F are referred to as Stueckelberg fields.

ii) Our Lagrangian for the free self-dual fields does not involve higher than second order terms in derivatives. Two-derivative contributions to the Lagrangian take the form of the standard kinetic terms of the antisymmetric tensor fields. The Lagrangian is invariant under gauge transformations and global conformal algebra transformations.

iii) Gauge transformations of the free self-dual fields do not involve higher than first order terms in derivatives. One-derivative contributions to the gauge transformations take the form of the standard gauge transformations of the antisymmetric tensor fields.

iv) The gauge symmetries of our Lagrangian make it possible to match our approach with the standard one, i.e., by an appropriate gauge fixing of the Stueckelberg fields and by solving

¹ It is self-duality of the field strength that leads to the problem with Lorentz invariant action for the gauge antisymmetric tensor field [2] without the use of auxiliary fields. Study of the Lorentz covariant formulations involving auxiliary fields may be found in Refs.[3, 4]. Interesting discussion of self-dual fields in $d = 6, 10$ may be found in [5].

²For instance, the self-dual field T^{ab} of $\mathcal{N} = 4, 4d$ conformal supergravity is described by the Lagrangian $\mathcal{L} = \partial^a \bar{T}^{ab} \partial^c T^{cb}$.

³ Making comparison with various approaches to massive fields, one can say that the standard approach to the self-dual fields is a counterpart of the Pauli-Fierz approach to the massive fields, while our approach to the self-dual fields is a counterpart of the Stueckelberg approach to the massive fields.

⁴ To realize those additional gauge symmetries we adopt the approach of Refs.[7, 8] which turns out to be the most useful for our purposes.

some constraints we obtain the standard formulation of the self-dual fields. This implies that our approach retain on-shell D.o.F of the standard theory of self-dual fields, i.e., on-shell, our approach is equivalent to the standard one.

As is well known, the Stueckelberg approach turned out to be successful for the study of theories involving massive fields. That is to say that all covariant formulations of string theories are realized by using Stueckelberg gauge symmetries. The self-dual fields enter field content of conformal supergravity theories. Therefore we expect that use of the Stueckelberg fields for the studying self-dual fields might be useful for developing new interesting formulations of the conformal supergravity theories.

The rest of the paper is organized as follows.

In Sec. 2, we summarize the notation and review the standard approach to the self-dual fields.

In Sec. 3, we start with the example of self-dual field propagating in $4d$ Minkowski space. For this field, we obtain the ordinary-derivative gauge invariant Lagrangian. We find realization of the conformal $so(4, 2)$ algebra symmetries on space of gauge fields and on space of field strengths. Also we obtain light-cone gauge Lagrangian and demonstrate that number of on-shell D.o.F of our approach coincides with the one in the standard approach to the self-dual field. We discuss the decomposition of those on-shell D.o.F into irreps of the $so(2)$ algebra.

In Sec. 4, we generalize results obtained in Sec. 3 to the case of self-dual fields propagating in Minkowski space of arbitrary dimension.

In Sec. 5, we represent our results in Secs. 3, 4 by using realization of field degrees of freedom in terms of generating functions. The generating functions are constructed out of the self-dual gauge fields and some oscillators. Use of the generating functions simplifies considerable study of the self-dual fields. Therefore we believe that result in Sec. 5 might be helpful in future studies of the self-dual fields.

Section 6 suggests directions for future research.

We collect various technical details in appendices. In Appendix A, we discuss details of the derivation of the ordinary-derivative gauge invariant Lagrangian. In Appendix B, we present details of the derivation of the conformal algebra transformations of gauge fields. In Appendix C, we discuss some details of the derivation of the light-cone gauge Lagrangian. In Appendix D we collect some useful formulas involving the Levi-Civita symbol.

2 Preliminaries

2.1 Notation

Throughout the paper, dimension of a flat space-time, which we denote by d , is restricted to be even integer, $d = 2\nu$. Coordinates in the flat space-time are denoted by x^a , while ∂_a stands for derivative with respect to x^a , $\partial_a \equiv \partial/\partial x^a$. Vector indices of the Lorentz algebra $so(d-1, 1)$ take the values $a, b, c, e = 0, 1, \dots, d-1$. To simplify our expressions we drop η_{ab} in scalar products, i.e. we use $X^a Y^a \equiv \eta_{ab} X^a Y^b$. The notation $\epsilon^{a_1 \dots a_\nu b_1 \dots b_\nu}$ stands for the Levi-Civita symbol. We assume the normalization $\epsilon^{01 \dots d-1} = 1$.

To avoid complicated tensor expressions we use a set of the creation operators $\alpha^a, \zeta, v^\oplus, v^\ominus$, and the respective set of annihilation operators $\bar{\alpha}^a, \bar{\zeta}, \bar{v}^\ominus, \bar{v}^\oplus$,

$$\bar{\alpha}^a|0\rangle = 0, \quad \bar{\zeta}|0\rangle = 0, \quad \bar{v}^\oplus|0\rangle = 0, \quad \bar{v}^\ominus|0\rangle = 0. \quad (2.1)$$

These operators satisfy the following (anti)commutation relations:

$$\{\bar{\alpha}^a, \alpha^b\} = \eta^{ab}, \quad \{\bar{\zeta}, \zeta\} = 1, \quad (2.2)$$

$$[\bar{v}^\oplus, v^\ominus] = 1, \quad [\bar{v}^\ominus, v^\oplus] = 1, \quad (2.3)$$

and will often be referred to as oscillators in what follows⁵. The oscillators $\alpha^a, \bar{\alpha}^a$ and $\zeta, \bar{\zeta}, v^\oplus, v^\ominus, \bar{v}^\oplus, \bar{v}^\ominus$ transform in the respective vector and scalar representations of the $so(d-1, 1)$ Lorentz algebra and satisfy the following hermitian conjugation rules:

$$\alpha^{a\dagger} = \bar{\alpha}^a, \quad \zeta^\dagger = \bar{\zeta}, \quad v^{\oplus\dagger} = \bar{v}^\oplus, \quad v^{\ominus\dagger} = \bar{v}^\ominus. \quad (2.4)$$

Throughout this paper we use operators constructed out of the oscillators and derivatives,

$$\square = \partial^a \partial_a, \quad \alpha \partial = \alpha^a \partial_a, \quad \bar{\alpha} \partial = \bar{\alpha}^a \partial_a, \quad (2.5)$$

$$N_\alpha \equiv \alpha^a \bar{\alpha}^a, \quad (2.6)$$

$$N_\zeta \equiv \zeta \bar{\zeta}, \quad (2.7)$$

$$N_{v^\oplus} \equiv v^\oplus \bar{v}^\oplus, \quad (2.8)$$

$$N_{v^\ominus} \equiv v^\ominus \bar{v}^\ominus, \quad (2.9)$$

$$N_v \equiv N_{v^\oplus} + N_{v^\ominus}. \quad (2.10)$$

2.2 Global conformal symmetries

In space-time of dimension d , the conformal algebra $so(d, 2)$ referred to the basis of Lorentz algebra $so(d-1, 1)$ consists of translation generators P^a , conformal boost generators K^a , dilatation generator D and generators $so(d-1, 1)$ Lorentz algebra J^{ab} . We assume the following normalization for commutators of the conformal algebra:

$$[D, P^a] = -P^a, \quad [P^a, J^{bc}] = \eta^{ab} P^c - \eta^{ac} P^b, \quad (2.11)$$

$$[D, K^a] = K^a, \quad [K^a, J^{bc}] = \eta^{ab} K^c - \eta^{ac} K^b, \quad (2.12)$$

$$[P^a, K^b] = \eta^{ab} D - J^{ab}, \quad (2.13)$$

$$[J^{ab}, J^{ce}] = \eta^{bc} J^{ae} + 3 \text{ terms}. \quad (2.14)$$

Let $|\phi\rangle$ denotes field propagating in flat space-time of dimension $d \geq 4$. Let Lagrangian for the free field $|\phi\rangle$ be conformal invariant. This implies, that Lagrangian is invariant with respect to transformation (invariance of the Lagrangian is assumed to be up to total derivative)

$$\delta_{\hat{G}} |\phi\rangle = \hat{G} |\phi\rangle, \quad (2.15)$$

⁵ We use oscillator formulation [9]-[14] to handle the many indices appearing for tensor fields. It can also be reformulated as an algebra acting on the symmetric-spinor bundle on the manifold M [15]. Note that the scalar oscillators $\zeta, \bar{\zeta}$ arise naturally by a dimensional reduction from flat space. It is natural to expect that the ‘conformal’ oscillators $v^\oplus, v^\ominus, \bar{v}^\oplus, \bar{v}^\ominus$ also allow certain interpretation via dimensional reduction. Interesting recent discussion of dimensional reduction may be found in Ref.[16].

where a realization of the conformal algebra generators $\hat{G}$ in terms of differential operators takes the form

$$P^a = \partial^a, \quad (2.16)$$

$$J^{ab} = x^a \partial^b - x^b \partial^a + M^{ab}, \quad (2.17)$$

$$D = x \partial + \Delta, \quad (2.18)$$

$$K^a = K_{\Delta, M}^a + R^a, \quad (2.19)$$

$$K_{\Delta, M}^a \equiv -\frac{1}{2} x^2 \partial^a + x^a D + M^{ab} x^b, \quad (2.20)$$

$$x \partial \equiv x^a \partial^a, \quad x^2 = x^a x^a. \quad (2.21)$$

In (2.17)-(2.19), Δ is operator of conformal dimension, M^{ab} is spin operator of the Lorentz algebra,

$$[M^{ab}, M^{ce}] = \eta^{bc} M^{ae} + 3 \text{ terms}, \quad (2.22)$$

and R^a is operator depending on the derivative ∂^a and not depending on the space-time coordinates x^a , $[P^a, R^b] = 0$. The spin operator M^{ab} is well known for arbitrary tensor fields of the Lorentz algebra. In the standard formulation of the self-dual fields, the operator R^a is equal to zero, while in the ordinary-derivative approach, we develop in this paper, the operator R^a is non-trivial. This implies that, in the framework of ordinary-derivative approach, the complete description of the self-dual fields requires finding not only gauge invariant Lagrangian but also the operator R^a .

Explicit representation for the action of operator $K_{\Delta, M}^a$ (2.20) is easily obtained from the relations above-given. Let $\Lambda^{a_1 \dots a_n}$ be rank- n antisymmetric tensor field of the Lorentz algebra $so(d-1, 1)$, while $\Delta(\Lambda)$ is a conformal dimension of this tensor field. Relation (2.19) implies that the conformal boost transformations of $\Lambda^{a_1 \dots a_n}$ can be presented as

$$\delta_{K^a} \Lambda^{a_1 \dots a_n} = \delta_{K_{\Delta, M}^a} \Lambda^{a_1 \dots a_n} + \delta_{R^a} \Lambda^{a_1 \dots a_n}, \quad (2.23)$$

$$\delta_{K_{\Delta, M}^a} \Lambda^{a_1 \dots a_n} = K_{\Delta(\Lambda)}^a \Lambda^{a_1 \dots a_n} + \sum_{k=1}^n M^{aa_k c} \Lambda^{a_1 \dots a_{k-1} c a_{k+1} \dots a_n}, \quad (2.24)$$

$$K_{\Delta}^a \equiv -\frac{1}{2} x^2 \partial^a + x^a (x \partial + \Delta), \quad (2.25)$$

$$M^{abc} \equiv \eta^{ab} x^c - \eta^{ac} x^b. \quad (2.26)$$

Thus, all that remains is to find explicit representation for the operator R^a . This is what we are doing, among other things, in this paper.

2.3 Standard approach to self-dual fields

We begin with brief review of the standard approach to the conformal self-dual fields. In this section we recall main facts of conformal field theory about these fields.

Consider totally antisymmetric rank- ν tensor field $T^{a_1 \dots a_\nu}$, $\nu \equiv \frac{d}{2}$, of the Lorentz algebra $so(d-1, 1)$, where the dimension of space-time d is even integer. In the framework of standard approach, the field $T^{a_1 \dots a_\nu}$ is referred to as conformal self-dual field if it meets the following requirements:

a) The field $T^{a_1 \dots a_\nu}$ satisfies the self-duality constraint

$$T^{a_1 \dots a_\nu} = \frac{\tau}{\nu!} \epsilon^{a_1 \dots a_\nu b_1 \dots b_\nu} T^{b_1 \dots b_\nu}, \quad (2.27)$$

$$\tau = \begin{cases} \pm i & \text{for } d = 4k; \\ \pm 1 & \text{for } d = 4k + 2; \end{cases} \quad (2.28)$$

where $\epsilon^{a_1 \dots a_\nu b_1 \dots b_\nu}$ is the Levi-Civita symbol. For flexibility, we do not fix sign of τ . Constraint (2.27) implies that $T^{a_1 \dots a_\nu}$ is *complex-valued* when $d = 4k$. In $d = 4k + 2$ the field $T^{a_1 \dots a_\nu}$ is considered to be *real-valued*.

b) Dynamics of the field $T^{a_1 \dots a_\nu}$ is described by the Lagrangian

$$\mathcal{L}_{\text{st}} = \frac{1}{(\nu - 1)!} \partial^a \bar{T}^{aa_2 \dots a_\nu} \partial^b T^{ba_2 \dots a_\nu}, \quad \text{for } d = 4k, \quad (2.29)$$

$$\mathcal{L}_{\text{st}} = \frac{1}{(\nu - 1)!} \partial^a T^{aa_2 \dots a_\nu} \partial^b T^{ba_2 \dots a_\nu}, \quad \text{for } d = 4k + 2, \quad (2.30)$$

where $\bar{T}$ in (2.29) stands for complex conjugate of T .

We now note that:

i) Requiring the Lagrangian to be invariant under the dilatation transformation we obtain the conformal dimension of the field $T^{a_1 \dots a_\nu}$,

$$\Delta(T^{a_1 \dots a_\nu}) = \frac{d - 2}{2}, \quad (2.31)$$

which is referred to as *the canonical conformal dimension of the conformal self-dual field*.

ii) Operator R^a of the field $T^{a_1 \dots a_\nu}$ is equal to zero.

iii) Simplest case of self-dual field, which is antisymmetric complex-valued rank-2 tensor field T^{ab} , corresponds to $d = 4$ with the following self-duality constraint (see (2.27))

$$T^{ab} = \frac{\tau}{2} \epsilon^{abce} T^{ce}. \quad (2.32)$$

The Lagrangian for the field T^{ab} can be read from (2.29)

$$\mathcal{L}_{\text{st}} = \partial^a \bar{T}^{ac} \partial^b T^{bc}. \quad (2.33)$$

The field T^{ab} appears in the field content of $\mathcal{N} = 4, 4d$ conformal supergravity.

3 Ordinary-derivative approach to self-dual field for $d = 4$

As a warm up let us start with the simplest case of the self-dual fields. Consider the self-dual field T^{ab} propagating in $4d$ flat space. In the framework of ordinary-derivative approach, a dynamical system that on-shell equivalent to the self-dual field T^{ab} with Lagrangian (2.33) involves two vector fields ϕ_{-1}^a and ϕ_1^a , one scalar field ϕ_0 and one self-dual rank-2 tensor field t^{ab} . In other words, we use the following field content:

$$\phi_{-1}^a, \quad \phi_1^a, \quad \phi_0, \quad t^{ab}, \quad (3.1)$$

$$t^{ab} = \frac{\tau}{2} \epsilon^{abce} t^{ce}, \quad (3.2)$$

$\tau = \pm i$. All fields in (3.1) are *complex-valued*. Conformal dimensions of these fields are given by

$$\Delta(\phi_{-1}^a) = 0, \quad \Delta(\phi_1^a) = 2, \quad \Delta(\phi_0) = 1, \quad \Delta(t^{ab}) = 1. \quad (3.3)$$

We note that subscript k' in $\phi_{k'}$ implies that conformal dimension of the field $\phi_{k'}$ is equal to $1 + k'$.

Ordinary-derivative action and Lagrangian we found take the form

$$S = \int d^4x \mathcal{L}, \quad (3.4)$$

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2} F^{ab}(\bar{\phi}_{-1}) F^{ab}(\phi_1) - \frac{1}{2} F^{ab}(\bar{\phi}_1) F^{ab}(\phi_{-1}) \\ & - \frac{1}{2} \bar{t}^{ab} F^{ab}(\phi_1) - \frac{1}{2} F^{ab}(\bar{\phi}_1) t^{ab} - (\bar{\phi}_1^a + \partial^a \bar{\phi}_0)(\phi_1^a + \partial^a \phi_0), \end{aligned} \quad (3.5)$$

where $\bar{\phi}_{k'}$ and $\bar{t}^{ab}$ are the respective complex conjugates of $\phi_{k'}$ and t^{ab} , while $F^{ab}(\phi)$ stands for field strength defined as

$$F^{ab}(\phi) \equiv \partial^a \phi^b - \partial^b \phi^a. \quad (3.6)$$

Details of the derivation of Lagrangian (3.5) may be found in Appendix A.

A few remarks are in order.

- i)** Two-derivative contributions to Lagrangian (3.5) are the standard kinetic terms for the vector fields ϕ_{-1}^a , ϕ_1^a and the standard Klein-Gordon kinetic term for the scalar field ϕ_0 .
- ii)** In addition to the two-derivative contributions, the Lagrangian involves one-derivative contributions and derivative-independent mass-like contributions. Appearance of the one-derivative and derivative-independent contributions to the Lagrangian is a characteristic feature of the ordinary-derivative approach.
- iii)** The self-dual field t^{ab} plays the role of a Lagrangian multiplier. Equations of motion for t^{ab} together with self-duality constraint for t^{ab} (3.2) tell us that *on-shell* the field strength $F^{ab}(\phi_1)$ is self-dual,

$$F^{ab}(\phi_1) = \frac{\tau}{2} \epsilon^{abce} F^{ce}(\phi_1). \quad (3.7)$$

Gauge transformations. To discuss gauge symmetries of Lagrangian (3.5) we introduce the following gauge transformation parameters:

$$\xi_{-2}, \quad \xi_0, \quad \lambda^a. \quad (3.8)$$

All these gauge transformation parameters are complex-valued. Conformal dimensions of the gauge transformation parameters are given by

$$\Delta(\xi_{-2}) = -1, \quad \Delta(\xi_0) = 1, \quad \Delta(\lambda^a) = 0. \quad (3.9)$$

We find that Lagrangian (3.5) is invariant under the gauge transformations

$$\delta \phi_1^a = \partial^a \xi_0, \quad (3.10)$$

$$\delta \phi_{-1}^a = \partial^a \xi_{-2} - \lambda^a, \quad (3.11)$$

$$\delta \phi_0 = -\xi_0, \quad (3.12)$$

$$\delta t^{ab} = F^{ab}(\lambda) + \frac{\tau}{2} \epsilon^{abce} F^{ce}(\lambda), \quad (3.13)$$

where field strength $F^{ab}(\lambda)$ for the gauge transformation parameter λ^a is defined as in (3.6).

From (3.11),(3.12), we see that the vector and scalar fields, ϕ_{-1}^a, ϕ_0 , transform as Stueckelberg fields, i.e., these fields can be gauged away via Stueckelberg gauge fixing, $\phi_{-1}^a = 0, \phi_0 = 0$. If we gauge away these fields, and exclude the vector field ϕ_1^a via equations of motion, then our Lagrangian reduces to the Lagrangian of standard approach (2.33). Note also that, in the Stueckelberg gauge, the field t^{ab} is identified with the generic self-dual field T^{ab} . Thus, our approach is equivalent to the standard one.

Realization of conformal algebra symmetries. To complete the ordinary-derivative description of the conformal self-dual field we should provide realization of the conformal algebra symmetries on a space of fields (3.1). The Poincaré algebra symmetries are realized on fields (3.1) in a standard way. Realization of dilatation symmetry is given by (2.18), where conformal dimensions of fields (3.1) are given in (3.3). General realization of conformal boost symmetries on arbitrary tensor fields is given in (2.23). According to (2.23) we should find realizations of the operators $K_{\Delta,M}^a$ and R^a on a space of fields (3.1). The realization of the former operator is obtained by adopting general formula (2.24) for gauge fields (3.1),

$$\delta_{K_{\Delta,M}^a} \phi_{k'}^b = K_{\Delta(\phi_{k'})}^a \phi_{k'}^b + M^{abc} \phi_{k'}^c, \quad k' = \pm 1, \quad (3.14)$$

$$\delta_{K_{\Delta,M}^a} \phi_0 = K_{\Delta(\phi_0)}^a \phi_0, \quad (3.15)$$

$$\delta_{K_{\Delta,M}^a} t^{a_1 a_2} = K_{\Delta(t)}^a t^{a_1 a_2} + M^{aa_1 c} t^{ca_2} + M^{aa_2 c} t^{a_1 c}, \quad (3.16)$$

where the operator K_{Δ}^a and conformal dimensions are defined in (2.25) and (3.3) respectively.

The real difficulty is to find the operator R^a . Realization of the operator R^a on a space of gauge fields (3.1) we found is given by

$$\delta_{R^a} \phi_1^b = -t^{ab} - \eta^{ab} \phi_0 - \partial^a \phi_{-1}^b - \frac{\tau}{2} \epsilon^{abce} F^{ce}(\phi_{-1}), \quad (3.17)$$

$$\delta_{R^a} \phi_{-1}^b = 0, \quad (3.18)$$

$$\delta_{R^a} \phi_0 = \phi_{-1}^a, \quad (3.19)$$

$$\delta_{R^a} t^{a_1 a_2} = \eta^{aa_1} \phi_{-1}^{a_2} - \eta^{aa_2} \phi_{-1}^{a_1} + \tau \epsilon^{a_1 a_2 ab} \phi_{-1}^b. \quad (3.20)$$

Using (3.14)-(3.20) and general formula (2.23),

$$\delta_{K^a} = \delta_{K_{\Delta,M}^a} + \delta_{R^a}, \quad (3.21)$$

gives the conformal boost transformations of the gauge fields.

From (3.17)-(3.20), we see the operator R^a maps the gauge field with conformal dimension Δ into the ones having conformal dimension less than Δ . This is to say that the realization of the operator R^a given in (3.17)-(3.20) can schematically be represented as

$$\phi_1 \xrightarrow{R} t \oplus \phi_0 \oplus \partial \phi_{-1}, \quad (3.22)$$

$$t \xrightarrow{R} \phi_{-1}, \quad \phi_0 \xrightarrow{R} \phi_{-1}, \quad \phi_{-1} \xrightarrow{R} 0. \quad (3.23)$$

Details of the derivation of the operator R^a may be found in Appendix B.⁶ As a side of remark we note that, having introduced field content and the Lagrangian, the operator R^a is fixed uniquely by requiring that

- i) the operator R^a should not involve higher than first order terms in the derivative;
- ii) Lagrangian should be invariant under the conformal algebra transformations.

As usually, the conformal algebra transformations of gauge fields (3.1) are defined up to gauge transformations. Alternatively, the conformal algebra symmetries can be realized on a space of field strengths. We now discuss field strengths for gauge fields (3.1) and the corresponding conformal transformations of the field strengths.

Realization of conformal algebra symmetries on space of field strengths. We introduce the following field strengths which are constructed out of gauge fields (3.1):

$$F^{\ominus ab} = T^{ab} , \quad (3.24)$$

$$F^{abc} = F^{abc}(T) , \quad (3.25)$$

$$F^{\oplus \ominus a} = \phi_1^a + \partial^a \phi_0 , \quad (3.26)$$

$$F^{\oplus ab} = F^{ab}(\phi_1) , \quad (3.27)$$

where $F^{ab}(\phi_1)$ is defined as in (3.6), while T^{ab} and $F^{abc}(T)$ are defined by the respective relations

$$T^{ab} = t^{ab} + F^{ab}(\phi_{-1}) + \frac{\tau}{2} \epsilon^{abce} F^{ce}(\phi_{-1}) , \quad (3.28)$$

$$F^{abc}(T) = \partial^a T^{bc} + \partial^b T^{ca} + \partial^c T^{ab} . \quad (3.29)$$

One can make sure that field strengths (3.24)-(3.27) are invariant under gauge transformations (3.10)-(3.13). Conformal dimensions of the field strengths can easily be read from relations (3.3) and (3.24)-(3.27)

$$\Delta(F^{\ominus ab}) = 1 , \quad \Delta(F^{abc}) = 2 , \quad \Delta(F^{\oplus \ominus a}) = 2 , \quad \Delta(F^{\oplus ab}) = 3 . \quad (3.30)$$

Poincaré algebra symmetries are realized on a space of the field strengths in a usual way. Realization of dilatation symmetry is given by (2.18), where conformal dimensions of the field strengths are given in (3.30). All that remains is to find conformal boost transformations of the field strengths. Making use of the conformal boost transformations of the gauge fields defined by relations (3.14)-(3.20), we find the corresponding conformal boost transformations of the field strengths. Those conformal boost transformations of the field strengths can be represented as in (2.23),(2.24), where we substitute, in place of the tensor fields $\Lambda^{a_1 \dots a_n}$, the fields strengths with the following realization of the operator R^a :

$$\delta_{R^a} F^{\ominus bc} = 0 , \quad (3.31)$$

$$\delta_{R^a} F^{bce} = -\eta^{ab} F^{\ominus ce} - \eta^{ac} F^{\ominus eb} - \eta^{ae} F^{\ominus bc} , \quad (3.32)$$

$$\delta_{R^a} F^{\oplus \ominus b} = -F^{\ominus ab} , \quad (3.33)$$

$$\delta_{R^a} F^{\oplus bc} = \eta^{ab} F^{\oplus \ominus c} - \eta^{ac} F^{\oplus \ominus b} + F^{abc} - \partial^a F^{\ominus bc} . \quad (3.34)$$

⁶ In appendix B, the operator R^a is obtained by using the realization of field D.o.F in terms of generating functions. Discussion of the generating functions may be found in Sec. 5. Therefore, before reading Appendix B, the reader should consult Sec. 5.

From (3.31)-(3.34), we see the operator R^a maps the field strength with conformal dimension Δ into the ones having conformal dimension less than Δ . In other words, the realization of the operator R^a given in (3.31)-(3.34) can schematically be represented as

$$F^\oplus \xrightarrow{R} F^{\oplus\ominus} \oplus F \oplus \partial F^\ominus, \quad (3.35)$$

$$F \xrightarrow{R} F^\ominus, \quad F^{\oplus\ominus} \xrightarrow{R} F^\ominus, \quad F^\ominus \xrightarrow{R} 0. \quad (3.36)$$

On-shell degrees of freedom and light-cone gauge Lagrangian. In order to discuss on-shell D.o.F of the conformal self-dual field we use a nomenclature of the $so(d-2)$ algebra which is $so(2)$ when $d=4$. Namely, we decompose the on-shell D.o.F into irreps of the $so(2)$ algebra. One can prove that the on-shell D.o.F of the self-dual field are described by two $so(2)$ algebra self-dual complex-valued vector fields ϕ_{-1}^i, ϕ_1^i and one complex-valued scalar fields ϕ_0 ,

$$\phi_{-1}^i, \quad \phi_1^i, \quad \phi_0, \quad (3.37)$$

where vector indices of the $so(2)$ algebra take values $i, j = 1, 2$. The vector fields satisfy the $so(2)$ self-duality constraint:

$$\phi_{-1}^i = \tau^{-1} \epsilon^{ij} \phi_{-1}^j, \quad \phi_1^i = \tau^{-1} \epsilon^{ij} \phi_1^j, \quad (3.38)$$

where ϵ^{ij} is the Levi-Civita symbol normalized as $\epsilon^{12} = 1$.

Using light-cone gauge frame, one can make sure that gauge invariant Lagrangian (3.5) leads to the following light-cone gauge Lagrangian for fields (3.37):

$$\mathcal{L}_{\text{l.c.}} = \bar{\phi}_1^i \square \phi_{-1}^i + \bar{\phi}_{-1}^i \square \phi_1^i + \bar{\phi}_0 \square \phi_0 - \bar{\phi}_1^i \phi_1^i. \quad (3.39)$$

Details of the derivation of the on-shell D.o.F and light-cone gauge Lagrangian may be found in Appendix C.

From (3.37),(3.38), we see that number of real-valued on-shell D.o.F is equal to 6. This result agrees with the one found in Ref.[6]. Note however that we find not only the number of the on-shell D.o.F but also provide the decomposition of those on-shell D.o.F into irreps of the $so(2)$ algebra.

4 Ordinary-derivative approach to self-dual field for arbitrary $d = 2\nu$

We now develop ordinary-derivative approach to conformal self-dual field propagating in flat space-time of arbitrary even dimension $d = 2\nu$. To discuss ordinary-derivative approach to the self-dual field we use the following field content:

$$\phi_{-1}^{a_1 \dots a_{\nu-1}}, \quad \phi_1^{a_1 \dots a_{\nu-1}}, \quad \phi_0^{a_1 \dots a_{\nu-2}}, \quad t^{a_1 \dots a_\nu}, \quad (4.1)$$

where the field $t^{a_1 \dots a_\nu}$ satisfies the self-duality constraint

$$t^{a_1 \dots a_\nu} = \frac{\tau}{\nu!} \epsilon^{a_1 \dots a_\nu b_1 \dots b_\nu} t^{b_1 \dots b_\nu}, \quad (4.2)$$

and τ is defined in (2.28). We note that:

- i) Fields in (4.1) are antisymmetric tensor fields of the Lorentz algebra $so(d-1, 1)$.
- ii) Fields in (4.1) are complex-valued when $d = 4k$ and real-valued when $d = 4k + 2$.
- iii) Conformal dimensions of fields in (4.1) are given by

$$\begin{aligned}\Delta(\phi_{-1}^{a_1 \dots a_{\nu-1}}) &= \frac{d-4}{2}, & \Delta(\phi_1^{a_1 \dots a_{\nu-1}}) &= \frac{d}{2}, & \Delta(\phi_0^{a_1 \dots a_{\nu-2}}) &= \frac{d-2}{2}, \\ \Delta(t^{a_1 \dots a_{\nu}}) &= \frac{d-2}{2}.\end{aligned}\quad (4.3)$$

We note that subscript k' in $\phi_{k'}$ implies that conformal dimension of the field $\phi_{k'}$ is equal to $\frac{d-2}{2} + k'$. Ordinary-derivative action we found is given by

$$S = \int d^d x \mathcal{L}, \quad (4.4)$$

where Lagrangian takes the form

$$\begin{aligned}\mathcal{L} &= -\frac{1}{\nu!} F^{a_1 \dots a_{\nu}}(\bar{\phi}_{-1}) F^{a_1 \dots a_{\nu}}(\phi_1) - \frac{1}{\nu!} F^{a_1 \dots a_{\nu}}(\bar{\phi}_1) F^{a_1 \dots a_{\nu}}(\phi_{-1}) \\ &- \frac{1}{\nu!} \bar{t}^{a_1 \dots a_{\nu}} F^{a_1 \dots a_{\nu}}(\phi_1) - \frac{1}{\nu!} t^{a_1 \dots a_{\nu}} F^{a_1 \dots a_{\nu}}(\bar{\phi}_1) \\ &- \frac{1}{(\nu-1)!} (\bar{\phi}_1^{a_1 \dots a_{\nu-1}} + F^{a_1 \dots a_{\nu-1}}(\bar{\phi}_0)) (\phi_1^{a_1 \dots a_{\nu-1}} + F^{a_1 \dots a_{\nu-1}}(\phi_0)),\end{aligned}\quad (4.5)$$

when $d = 4k$, while for $d = 4k + 2$, Lagrangian is given by

$$\begin{aligned}\mathcal{L} &= -\frac{1}{\nu!} F^{a_1 \dots a_{\nu}}(\phi_{-1}) F^{a_1 \dots a_{\nu}}(\phi_1) - \frac{1}{\nu!} t^{a_1 \dots a_{\nu}} F^{a_1 \dots a_{\nu}}(\phi_1) \\ &- \frac{1}{2(\nu-1)!} (\phi_1^{a_1 \dots a_{\nu-1}} + F^{a_1 \dots a_{\nu-1}}(\phi_0)) (\phi_1^{a_1 \dots a_{\nu-1}} + F^{a_1 \dots a_{\nu-1}}(\phi_0)),\end{aligned}\quad (4.6)$$

where field strengths are defined as

$$F^{a_1 \dots a_n}(\phi) = n \partial^{[a_1} \phi^{a_2 \dots a_n]} \quad (4.7)$$

and the antisymmetrization of the tensor indices is normalized as $[a_1 \dots a_n] = \frac{1}{n!} (a_1 \dots a_n \pm (n-1) \text{ terms})$.

We note that:

- i) Two-derivative contributions to Lagrangians (4.5),(4.6) take the form of standard second-order kinetic terms for antisymmetric tensor fields. Besides the two-derivative contributions, the Lagrangians involve one-derivative contributions and derivative-independent mass-like contributions.
- ii) Equations of motion for $t^{a_1 \dots a_{\nu}}$ and self-duality constraint for $t^{a_1 \dots a_{\nu}}$ (4.2) imply that *on-shell* the field strength $F^{a_1 \dots a_{\nu}}(\phi_1)$ satisfies the self-duality constraint,

$$F^{a_1 \dots a_{\nu}}(\phi_1) = \frac{\tau}{\nu!} \epsilon^{a_1 \dots a_{\nu} b_1 \dots b_{\nu}} F^{b_1 \dots b_{\nu}}(\phi_1). \quad (4.8)$$

Gauge transformations. We now discuss gauge symmetries of Lagrangians (4.5),(4.6). To this end we introduce the following gauge transformation parameters:

$$\xi_{-2}^{a_1 \dots a_{\nu-2}}, \quad \xi_0^{a_1 \dots a_{\nu-2}}, \quad \xi_{-1}^{a_1 \dots a_{\nu-3}}, \quad \lambda^{a_1 \dots a_{\nu-1}}. \quad (4.9)$$

We note that:

- i) Gauge transformation parameters (4.9) are antisymmetric tensor fields of the Lorentz algebra

$so(d-1, 1)$.

ii) Gauge transformation parameters (4.9) are complex-valued when $d = 4k$ and real-valued when $d = 4k + 2$.

iii) Conformal dimensions of gauge transformation parameters (4.9) are given by

$$\begin{aligned}\Delta(\xi_{-1}^{a_1 \dots a_{\nu-2}}) &= \frac{d-6}{2}, & \Delta(\xi_0^{a_1 \dots a_{\nu-2}}) &= \frac{d-2}{2}, & \Delta(\xi_{-1}^{a_1 \dots a_{\nu-3}}) &= \frac{d-4}{2}, \\ \Delta(\lambda^{a_1 \dots a_{\nu-1}}) &= \frac{d-4}{2}.\end{aligned}\tag{4.10}$$

Gauge transformations we found take the form:

$$\delta\phi_1^{a_1 \dots a_{\nu-1}} = F^{a_1 \dots a_{\nu-1}}(\xi_0), \tag{4.11}$$

$$\delta\phi_{-1}^{a_1 \dots a_{\nu-1}} = F^{a_1 \dots a_{\nu-1}}(\xi_{-2}) - \lambda^{a_1 \dots a_{\nu-1}}, \tag{4.12}$$

$$\delta\phi_0^{a_1 \dots a_{\nu-2}} = F^{a_1 \dots a_{\nu-2}}(\xi_{-1}) - \xi_0^{a_1 \dots a_{\nu-2}}, \tag{4.13}$$

$$\delta t^{a_1 \dots a_{\nu}} = F^{a_1 \dots a_{\nu}}(\lambda) + \frac{\tau}{\nu!} \epsilon^{a_1 \dots a_{\nu} b_1 \dots b_{\nu}} F^{b_1 \dots b_{\nu}}(\lambda), \tag{4.14}$$

where strengths for the gauge transformation parameters are defined as in (4.7).

From (4.12),(4.13), we see that the gauge fields $\phi_{-1}^{a_1 \dots a_{\nu-1}}$, $\phi_0^{a_1 \dots a_{\nu-2}}$ transform as Stueckelberg fields, i.e., these fields can be gauged away via Stueckelberg gauge fixing, $\phi_{-1}^{a_1 \dots a_{\nu-1}} = 0$, $\phi_0^{a_1 \dots a_{\nu-2}} = 0$. If we gauge away these fields and exclude the field $\phi_1^{a_1 \dots a_{\nu-1}}$ via equations of motion, then our Lagrangians (4.5),(4.6) reduce to the respective Lagrangians of the standard approach, (2.29),(2.30). Thus, our approach is equivalent to the standard one. Note that one-derivative contributions to gauge transformations (4.11)-(4.14) take the form of the standard gauge transformations for antisymmetric tensor fields.

Realization of conformal algebra symmetries. To complete the ordinary-derivative description of the conformal self-dual field we should provide realization of the conformal algebra symmetries on a space of tensor fields (4.1). Realization of the Poincaré algebra symmetries on the tensor fields is well known. Realization of dilatation symmetry is given by (2.18), where conformal dimensions of fields (4.1) are given in (4.3). General form of conformal boost transformations of arbitrary tensor fields is given in (2.23). According to (2.23) we should find realizations of the operators $K_{\Delta, M}^a$ and R^a on space of fields (4.1). The realization of the former operator is obtained by adopting general formula (2.24) for gauge fields (4.1). The real problem is getting a realization of the operator R^a . The realization of the operator R^a on space of gauge fields (4.1) we found is given by

$$\begin{aligned}\delta_{R^a} \phi_1^{a_1 \dots a_{\nu-1}} &= -t^{aa_1 \dots a_{\nu-1}} - (\nu-1)\eta^{a[a_1} \phi_0^{a_2 \dots a_{\nu-1}]} \\ &- \partial^a \phi_{-1}^{a_1 \dots a_{\nu-1}} - \frac{\tau}{\nu!} \epsilon^{aa_1 \dots a_{\nu-1} b_1 \dots b_{\nu}} F^{b_1 \dots b_{\nu}}(\phi_{-1}),\end{aligned}\tag{4.15}$$

$$\delta_{R^a} \phi_{-1}^{a_1 \dots a_{\nu-1}} = 0, \tag{4.16}$$

$$\delta_{R^a} \phi_0^{a_1 \dots a_{\nu-2}} = \phi_{-1}^{aa_1 \dots a_{\nu-2}}, \tag{4.17}$$

$$\delta_{R^a} t^{a_1 \dots a_{\nu}} = \nu \eta^{a[a_1} \phi_{-1}^{a_2 \dots a_{\nu}]} + \frac{\tau}{(\nu-1)!} \epsilon^{a_1 \dots a_{\nu} ab_1 \dots b_{\nu-1}} \phi_{-1}^{b_1 \dots b_{\nu-1}}. \tag{4.18}$$

Making use of these relations and general formula (2.23) gives the conformal boost transformations of gauge fields (4.1).

On-shell degrees of freedom and light-cone gauge Lagrangian. We now discuss on-shell D.o.F of the self-dual field. As before, for this purpose it is convenient to use fields transforming in irreps of the $so(d-2)$ algebra. Using the method in Appendix C, one can prove that on-shell D.o.F are described by the following antisymmetric tensor fields of the $so(d-2)$ algebra:

$$\phi_{-1}^{i_1 \dots i_{\nu-1}}, \quad \phi_1^{i_1 \dots i_{\nu-1}}, \quad \phi_0^{i_1 \dots i_{\nu-2}}, \quad (4.19)$$

where vector indices of the $so(d-2)$ algebra take values $i, j = 1, 2, \dots, d-2$. Fields in (4.19) are complex-valued when $d = 4k$ and real-valued when $d = 4k + 2$. The fields $\phi_{-1}^{i_1 \dots i_{\nu-1}}, \phi_1^{i_1 \dots i_{\nu-1}}$ satisfy the $so(d-2)$ self-duality constraint,

$$\phi_{-1}^{i_1 \dots i_{\nu-1}} = \frac{\tau^{-1}}{(\nu-1)!} \epsilon^{i_1 \dots i_{\nu-1} j_1 \dots j_{\nu-1}} \phi_{-1}^{j_1 \dots j_{\nu-1}}, \quad (4.20)$$

$$\phi_1^{i_1 \dots i_{\nu-1}} = \frac{\tau^{-1}}{(\nu-1)!} \epsilon^{i_1 \dots i_{\nu-1} j_1 \dots j_{\nu-1}} \phi_1^{j_1 \dots j_{\nu-1}}, \quad (4.21)$$

where $\epsilon^{i_1 \dots i_{\nu-1} j_1 \dots j_{\nu-1}}$ is the Levi-Civita symbol normalized as $\epsilon^{12 \dots d-2} = 1$.

Total number of real-valued on-shell D.o.F given in (4.19) is equal to

$$\mathbf{n} = \frac{h(5\nu-7)(2\nu-4)!}{(\nu-1)!(\nu-2)!}, \quad h = \begin{cases} 2 & \text{for } d = 4k; \\ 1 & \text{for } d = 4k + 2. \end{cases} \quad (4.22)$$

Namely, we note that $\mathbf{n}$ is a sum of $\mathbf{n}(\phi_{\pm 1}^{i_1 \dots i_{\nu-1}})$ and $\mathbf{n}(\phi_0^{i_1 \dots i_{\nu-2}})$ which are the respective numbers of the real-valued independent tensorial components of the fields $\phi_{\pm 1}^{i_1 \dots i_{\nu-1}}$ and $\phi_0^{i_1 \dots i_{\nu-2}}$,

$$\mathbf{n} = \mathbf{n}(\phi_1^{i_1 \dots i_{\nu-1}}) + \mathbf{n}(\phi_{-1}^{i_1 \dots i_{\nu-1}}) + \mathbf{n}(\phi_0^{i_1 \dots i_{\nu-2}}), \quad (4.23)$$

$$\mathbf{n}(\phi_{\pm 1}^{i_1 \dots i_{\nu-1}}) = \frac{h(2\nu-2)!}{2((\nu-1)!)^2}, \quad \mathbf{n}(\phi_0^{i_1 \dots i_{\nu-2}}) = \frac{h(2\nu-4)!}{((\nu-2)!)^2}. \quad (4.24)$$

Using light-cone gauge frame, one can make sure that, for $d = 4k$, gauge invariant Lagrangian (4.5) leads to the following light-cone gauge Lagrangian for fields (4.19):

$$\begin{aligned} \mathcal{L}_{\text{l.c.}} = & \frac{1}{(\nu-1)!} \bar{\phi}_1^{i_1 \dots i_{\nu-1}} \square \phi_{-1}^{i_1 \dots i_{\nu-1}} + \frac{1}{(\nu-1)!} \bar{\phi}_{-1}^{i_1 \dots i_{\nu-1}} \square \phi_1^{i_1 \dots i_{\nu-1}} \\ & + \frac{1}{(\nu-2)!} \bar{\phi}_0^{i_1 \dots i_{\nu-2}} \square \phi_0^{i_1 \dots i_{\nu-2}} - \frac{1}{(\nu-1)!} \bar{\phi}_1^{i_1 \dots i_{\nu-1}} \phi_1^{i_1 \dots i_{\nu-1}}, \end{aligned} \quad (4.25)$$

while, for $d = 4k + 2$, gauge invariant Lagrangian (4.6) leads to the following light-cone gauge Lagrangian for fields (4.19):

$$\begin{aligned} \mathcal{L}_{\text{l.c.}} = & \frac{1}{(\nu-1)!} \phi_1^{i_1 \dots i_{\nu-1}} \square \phi_{-1}^{i_1 \dots i_{\nu-1}} + \frac{1}{2(\nu-2)!} \phi_0^{i_1 \dots i_{\nu-2}} \square \phi_0^{i_1 \dots i_{\nu-2}} \\ & - \frac{1}{2(\nu-1)!} \phi_1^{i_1 \dots i_{\nu-1}} \phi_1^{i_1 \dots i_{\nu-1}}. \end{aligned} \quad (4.26)$$

5 Oscillator form of Lagrangian

In the preceding sections, we have presented our results for the self-dual fields by using the representation of field content in terms of the tensor fields. However use of such representation is not convenient in many applications. In this section, we represent our results by using the representation of field content in terms of generating functions constructed out of the tensor fields and the appropriate oscillators.⁷ This is to say that in order to obtain the Lagrangian description in an easy-to-use form, we introduce creation operators α^a , ζ , $v^\oplus$, $v^\ominus$ and the respective annihilation operators $\bar{\alpha}^a$, $\bar{\zeta}$, $\bar{v}^\ominus$, $\bar{v}^\oplus$ and collect tensor fields (4.1) in ket-vector $|\Phi\rangle$ defined by

$$|\Phi\rangle = \begin{pmatrix} |\phi\rangle \\ |t\rangle \end{pmatrix}, \quad (5.1)$$

$$|\phi\rangle = v^\oplus|\phi_1\rangle + v^\ominus|\phi_{-1}\rangle + \zeta|\phi_0\rangle, \quad (5.2)$$

$$|\phi_{k'}\rangle \equiv \frac{1}{(\nu-1)!} \alpha^{a_1} \dots \alpha^{a_{\nu-1}} \phi_{k'}^{a_1 \dots a_{\nu-1}} |0\rangle, \quad k' = -1, 1, \quad (5.3)$$

$$|\phi_0\rangle \equiv \frac{1}{(\nu-2)!} \alpha^{a_1} \dots \alpha^{a_{\nu-2}} \phi_0^{a_1 \dots a_{\nu-2}} |0\rangle, \quad (5.4)$$

$$|t\rangle \equiv \frac{1}{\nu!} \alpha^{a_1} \dots \alpha^{a_\nu} t^{a_1 \dots a_\nu} |0\rangle. \quad (5.5)$$

In the literature, ket-vectors (5.1)-(5.5) are sometimes referred to as generating functions. The ket-vectors $|\phi\rangle$, $|t\rangle$ satisfy the obvious algebraic constraints

$$(N_\alpha + N_\zeta)|\phi\rangle = (\nu-1)|\phi\rangle, \quad (5.6)$$

$$(N_\zeta + N_v)|\phi\rangle = |\phi\rangle, \quad (5.7)$$

$$N_\alpha|t\rangle = \nu|t\rangle, \quad (5.8)$$

$$N_\zeta|t\rangle = 0, \quad N_v|t\rangle = 0, \quad (5.9)$$

where we use the notation given in (2.6)-(2.10). We note that these algebraic constraints tell us about number of the oscillators α^a , ζ , $v^\oplus$, $v^\ominus$ appearing in the ket-vectors $|\phi\rangle$ and $|t\rangle$. In terms of the ket-vector $|t\rangle$, self-duality constraint (4.2) takes the form

$$|t\rangle = \epsilon|t\rangle, \quad (5.10)$$

where we use the notation for ϵ -symbol

$$\epsilon \equiv \frac{\tau}{(\nu!)^2} \epsilon^{a_1 \dots a_\nu b_1 \dots b_\nu} \alpha^{a_1} \dots \alpha^{a_\nu} \bar{\alpha}^{b_\nu} \dots \bar{\alpha}^{b_1}. \quad (5.11)$$

Useful relations for ϵ -symbol (5.11) and various related ϵ -symbols may be found in Appendix D.

⁷ Note that in this paper we use oscillators just to handle the many indices appearing for tensor fields. In a proper way, the oscillators arise in the framework of world-line approach to higher-spin fields (see e.g. Refs.[17]-[19].)

In terms of the ket-vector $|\Phi\rangle$, Lagrangians (4.5),(4.6) can be reexpressed as

$$\mathcal{L} = \frac{h}{2} \langle \Phi | E | \Phi \rangle , \quad (5.12)$$

where the normalization factor h is given in (4.22), while operator E is defined by the relations

$$E = \begin{pmatrix} E_{\phi\phi} & E_{\phi t} \\ E_{t\phi} & 0 \end{pmatrix} , \quad (5.13)$$

$$E_{\phi\phi} = E_{\phi\phi(2)} + E_{\phi\phi(1)} + E_{\phi\phi(0)} , \quad (5.14)$$

$$E_{\phi t} = v^\ominus \bar{\alpha} \partial , \quad E_{t\phi} = -\bar{v}^\ominus \alpha \partial , \quad (5.15)$$

$$E_{\phi\phi(2)} = \square - \alpha \partial \bar{\alpha} \partial , \quad (5.16)$$

$$E_{\phi\phi(1)} = e_1 \bar{\alpha} \partial + \alpha \partial \bar{e}_1 , \quad (5.17)$$

$$E_{\phi\phi(0)} = m_1 , \quad (5.18)$$

$$e_1 = \zeta \bar{v}^\ominus , \quad \bar{e}_1 = -v^\ominus \bar{\zeta} , \quad m_1 = v^\ominus \bar{v}^\ominus (N_\zeta - 1) . \quad (5.19)$$

Alternatively, Lagrangian (5.12) can be represented in terms of ket-vector of gauge fields (5.2)-(5.5). This is to say that the Lagrangian takes the form (up to a total derivative)

$$\begin{aligned} -\mathcal{L} &= \langle F(\phi_{-1}) | F(\phi_1) \rangle + \langle F(\phi_1) | F(\phi_{-1}) \rangle + \langle F(\phi_1) | t \rangle + \langle t | F(\phi_1) \rangle \\ &+ \left(\langle \phi_1 | + \langle F(\phi_0) | \right) \left(| \phi_1 \rangle + | F(\phi_0) \rangle \right) , \end{aligned} \quad (5.20)$$

when $d = 4k$, and

$$\begin{aligned} -\mathcal{L} &= \langle F(\phi_{-1}) | F(\phi_1) \rangle + \langle F(\phi_1) | t \rangle \\ &+ \frac{1}{2} \left(\langle \phi_1 | + \langle F(\phi_0) | \right) \left(| \phi_1 \rangle + | F(\phi_0) \rangle \right) , \end{aligned} \quad (5.21)$$

when $d = 4k + 2$. In (5.20), (5.21) and below, ket-vector of field strength $|F(\phi)\rangle$ is defined as

$$|F(\phi)\rangle = \alpha \partial | \phi \rangle . \quad (5.22)$$

Gauge transformations. Gauge transformations can also be cast into the generating form. To this end we use, as before, the oscillators $\alpha^a, \zeta, v^\oplus, v^\ominus$ and collect gauge transformation parameters (4.9) into ket-vector $|\Xi\rangle$ defined by

$$|\Xi\rangle = \begin{pmatrix} |\xi\rangle \\ |\lambda\rangle \end{pmatrix} , \quad (5.23)$$

where ket-vectors $|\xi\rangle$ and $|\lambda\rangle$ are defined as

$$|\xi\rangle = v^\oplus |\xi_0\rangle + v^\ominus |\xi_{-2}\rangle - \zeta |\xi_{-1}\rangle, \quad (5.24)$$

$$|\xi_{k'-1}\rangle \equiv \frac{1}{(\nu-2)!} \alpha^{a_1} \dots \alpha^{a_{\nu-2}} \xi_{k'-1}^{a_1 \dots a_{\nu-2}} |0\rangle, \quad k' = -1, 1, \quad (5.25)$$

$$|\xi_{-1}\rangle \equiv \frac{1}{(\nu-3)!} \alpha^{a_1} \dots \alpha^{a_{\nu-3}} \xi_{-1}^{a_1 \dots a_{\nu-3}} |0\rangle, \quad (5.26)$$

$$|\lambda\rangle \equiv \frac{1}{(\nu-1)!} \alpha^{a_1} \dots \alpha^{a_{\nu-1}} \lambda^{a_1 \dots a_{\nu-1}} |0\rangle. \quad (5.27)$$

The ket-vectors $|\xi\rangle, |\lambda\rangle$ satisfy the obvious algebraic constraints

$$(N_\alpha + N_\zeta) |\xi\rangle = (\nu-2) |\xi\rangle, \quad (5.28)$$

$$(N_\zeta + N_v) |\xi\rangle = |\xi\rangle, \quad (5.29)$$

$$N_\alpha |\lambda\rangle = (\nu-1) |\lambda\rangle, \quad (5.30)$$

$$N_\zeta |\lambda\rangle = 0, \quad N_v |\lambda\rangle = 0. \quad (5.31)$$

As before, these constraints tell us about number of the oscillators $\alpha^a, \zeta, v^\oplus, v^\ominus$ appearing in the ket-vectors $|\xi\rangle$ and $|\lambda\rangle$.

Now, gauge transformations (4.11)-(4.14) can entirely be represented in terms of the ket-vectors $|\Phi\rangle$ and $|\Xi\rangle$,

$$\delta |\Phi\rangle = G |\Xi\rangle, \quad (5.32)$$

where the operator G is given by

$$G = \begin{pmatrix} \alpha \partial - \zeta \bar{v}^\ominus & -v^\ominus \\ 0 & (1 + \epsilon) \alpha \partial \end{pmatrix}, \quad (5.33)$$

and the ϵ -symbol is defined in (5.11).

Alternatively, gauge transformation (5.32) can be represented in terms of ket-vector of gauge fields (5.2)-(5.5). This is to say that gauge transformation (5.32) amounts to the following gauge transformations

$$\delta |\phi_1\rangle = |F(\xi_0)\rangle, \quad (5.34)$$

$$\delta |\phi_{-1}\rangle = |F(\xi_{-2})\rangle - |\lambda\rangle, \quad (5.35)$$

$$\delta |\phi_0\rangle = |F(\xi_{-1})\rangle - |\xi_0\rangle, \quad (5.36)$$

$$\delta |t\rangle = (1 + \epsilon) |F(\lambda)\rangle, \quad (5.37)$$

where the field strengths for ket-vectors of the gauge transformation parameters are defined as in (5.22).

Oscillator realization of conformal algebra symmetries on gauge fields. To complete the oscillator description of the conformal self-dual field we provide a realization of the conformal algebra symmetries on space of the ket-vector $|\Phi\rangle$. Realization of the Poincaré algebra symmetries and dilatation symmetry is given by (2.16)-(2.18), where the operators M^{ab} and Δ take the form

$$M^{ab} = \alpha^a \bar{\alpha}^b - \alpha^b \bar{\alpha}^a, \quad (5.38)$$

$$\Delta = \frac{d-2}{2} + \Delta', \quad \Delta' \equiv N_{v^\oplus} - N_{v^\ominus}. \quad (5.39)$$

Conformal boost transformations of $|\Phi\rangle$ are given in (2.19). According to (2.19) we should find the operators $K_{\Delta,M}^a$ and R^a . The former operator is given in (2.20). Realization of the operator R^a on space of $|\Phi\rangle$ can be read from (4.15)-(4.18). Namely, in terms of the ket-vector $|\Phi\rangle$, transformations of the gauge fields given in (4.15)-(4.18) can be represented as

$$\delta_{R^a} |\Phi\rangle = R^a |\Phi\rangle, \quad (5.40)$$

where the realization of the operator R^a on $|\Phi\rangle$ takes the form

$$R^a = \begin{pmatrix} R_{\phi\phi}^a & R_{\phi t}^a \\ R_{t\phi}^a & 0 \end{pmatrix}, \quad (5.41)$$

$$R_{\phi\phi}^a = r_{0,1} \bar{\alpha}^a + \alpha^a \bar{r}_{0,1} + r_{1,1} (\eta^{ab} + \epsilon_0^{ab}) \partial^b, \quad (5.42)$$

$$R_{\phi t}^a = r_{0,4} \bar{\alpha}^a, \quad (5.43)$$

$$R_{t\phi}^a = r_{0,5} (1 + \epsilon) \alpha^a, \quad (5.44)$$

$$r_{0,1} = \zeta \bar{v}^\oplus, \quad \bar{r}_{0,1} = -v^\oplus \bar{\zeta}, \quad (5.45)$$

$$r_{1,1} = -v^\oplus \bar{v}^\oplus, \quad (5.46)$$

$$r_{0,4} = -v^\oplus, \quad r_{0,5} = \bar{v}^\oplus, \quad (5.47)$$

and ϵ_0^{ab} -symbol is defined in (D.4).

Alternatively, conformal boost transformations (5.40) can be represented in terms of ket-vectors (5.2)-(5.5). To this end we note that realization of the operator $K_{\Delta,M}^a$ on space of ket-vectors (5.2)-(5.5) is given by (2.20), where the operators M^{ab} and Δ take the same form as in (5.38),(5.39). Note that (5.39) implies the following conformal dimensions of the respective ket-vectors

$$\Delta(|\phi_{k'}\rangle) = \frac{d-2}{2} + k', \quad k' = 0, \pm 1, \quad (5.48)$$

$$\Delta(|t\rangle) = \frac{d-2}{2}. \quad (5.49)$$

We now make sure that the realization of the operator R^a given in (5.40) can be represented in terms of ket-vectors (5.2)-(5.5) as

$$\delta_{R^a} |\phi_1\rangle = -\bar{\alpha}^a |t\rangle - \alpha^a |\phi_0\rangle - (\eta^{ab} + \epsilon_0^{ab}) \partial^b |\phi_{-1}\rangle, \quad (5.50)$$

$$\delta_{R^a} |\phi_{-1}\rangle = 0, \quad (5.51)$$

$$\delta_{R^a} |\phi_0\rangle = \bar{\alpha}^a |\phi_{-1}\rangle, \quad (5.52)$$

$$\delta_{R^a} |t\rangle = (1 + \epsilon) \alpha^a |\phi_{-1}\rangle. \quad (5.53)$$

Making use of these relations gives the conformal boost transformations of ket-vectors (5.2)-(5.5).

We now remind that the realization of the conformal symmetries on a space of the ket-vectors of gauge fields (5.50)-(5.53) is defined up to gauge transformations (5.34)-(5.37). As we have demonstrated in Sec. 3, the conformal symmetries can also be realized on a space of the field strengths. We now discuss the oscillator form of the field strengths for gauge fields (5.2)-(5.5) and the corresponding realization of conformal symmetries on a space of the field strengths.

Oscillator realization of conformal algebra symmetries on field strengths. We introduce the following ket-vectors of field strengths constructed out of ket-vectors of gauge fields (5.2)-(5.5):

$$|F^\ominus\rangle = |t\rangle + (1 + \epsilon)|F(\phi_{-1})\rangle, \quad (5.54)$$

$$|F\rangle = \alpha\partial|F^\ominus\rangle, \quad (5.55)$$

$$|F^{\oplus\ominus}\rangle = |\phi_1\rangle + |F(\phi_0)\rangle, \quad (5.56)$$

$$|F^\oplus\rangle = |F(\phi_1)\rangle, \quad (5.57)$$

where $|F(\phi_{k'})\rangle$, $k' = 0, \pm 1$ are defined as in (5.22). One can make sure that field strengths (5.54)-(5.57) are invariant under gauge transformations (5.34)-(5.37).

Conformal dimensions of the field strengths can be read from (5.48),(5.49) and (5.54)-(5.57),

$$\Delta(|F^\ominus\rangle) = \frac{d-2}{2}, \quad \Delta(|F\rangle) = \frac{d}{2}, \quad \Delta(|F^{\oplus\ominus}\rangle) = \frac{d}{2}, \quad \Delta(|F^\oplus\rangle) = \frac{d+2}{2}. \quad (5.58)$$

Realization of the Poincaré algebra symmetries and dilatation symmetry on space of the ket-vectors of field strengths is given by (2.16)-(2.18), where the operator M^{ab} takes the form as in (5.38), while conformal dimensions are given in (5.58).

Making use of transformations of the ket-vectors of gauge fields given in (5.50)-(5.53), we find the corresponding conformal boost transformations of the ket-vectors of field strengths,

$$\delta_{K^a}|F^\ominus\rangle = K_{\Delta,M}^a|F^\ominus\rangle, \quad (5.59)$$

$$\delta_{K^a}|F\rangle = K_{\Delta,M}^a|F\rangle - \alpha^a|F^\ominus\rangle, \quad (5.60)$$

$$\delta_{K^a}|F^{\oplus\ominus}\rangle = K_{\Delta,M}^a|F^{\oplus\ominus}\rangle - \bar{\alpha}^a|F^\ominus\rangle, \quad (5.61)$$

$$\delta_{K^a}|F^\oplus\rangle = K_{\Delta,M}^a|F^\oplus\rangle + \alpha^a|F^{\oplus\ominus}\rangle + \bar{\alpha}^a|F\rangle - \partial^a|F^\ominus\rangle, \quad (5.62)$$

where the operator $K_{\Delta,M}^a$ is given in (2.20), while the conformal dimensions are defined in (5.58). Comparing these formulas with general relation (2.19), we find the realization of the operator R^a on a space of the ket-vectors of the field strengths,

$$\delta_{R^a}|F^\ominus\rangle = 0, \quad (5.63)$$

$$\delta_{R^a}|F\rangle = -\alpha^a|F^\ominus\rangle, \quad (5.64)$$

$$\delta_{R^a}|F^{\oplus\ominus}\rangle = -\bar{\alpha}^a|F^\ominus\rangle, \quad (5.65)$$

$$\delta_{R^a}|F^\oplus\rangle = \alpha^a|F^{\oplus\ominus}\rangle + \bar{\alpha}^a|F\rangle - \partial^a|F^\ominus\rangle. \quad (5.66)$$

Oscillator form of the light-cone gauge Lagrangian. To discuss oscillator form of light-cone gauge Lagrangian we collect fields (4.19) into the following ket-vectors:

$$|\phi_{1.c.}\rangle = v^{\oplus}|\phi_1\rangle_{1.c} + v^{\ominus}|\phi_{-1}\rangle_{1.c} + \zeta|\phi_0\rangle_{1.c}, \quad (5.67)$$

$$|\phi_{k'}\rangle_{1.c} \equiv \frac{1}{(\nu-1)!} \alpha^{i_1} \dots \alpha^{i_{\nu-1}} \phi_{k'}^{i_1 \dots i_{\nu-1}} |0\rangle, \quad k' = -1, 1, \quad (5.68)$$

$$|\phi_0\rangle_{1.c} \equiv \frac{1}{(\nu-2)!} \alpha^{i_1} \dots \alpha^{i_{\nu-2}} \phi_0^{i_1 \dots i_{\nu-2}} |0\rangle. \quad (5.69)$$

In terms of the ket-vector $|\phi_{1.c.}\rangle$ (5.67), light-cone gauge Lagrangians (4.25),(4.26) can concisely be represented as

$$\mathcal{L}_{1.c.} = \frac{h}{2} \langle \phi_{1.c.} | (\square - \mathcal{M}^2) | \phi_{1.c.} \rangle, \quad (5.70)$$

$$\mathcal{M}^2 \equiv v^{\ominus} \bar{v}^{\ominus}, \quad (5.71)$$

where the normalization factor h is given in (4.22).

6 Conclusions

In this paper, we applied the ordinary-derivative approach, developed in Ref.[25], to the study of conformal self-dual fields in flat space of even dimension. The results presented here should have a number of interesting applications and generalizations, some of which are:

i) Results in this paper and the ones in Ref.[25] provide the complete ordinary-derivative description of all fields that appear in the graviton supermultiplets of conformal supergravity theories. It would be interesting to apply these results to the study of supersymmetric conformal field theories [20]-[24] in the framework of ordinary-derivative approach. The first step in this direction would be understanding of how the supersymmetries are realized in the framework of our approach.

ii) Our approach to conformal theories (see Refs.[25, 26]) is based on the new realization of conformal gauge symmetries via Stueckelberg fields. In our approach, use of the Stueckelberg fields is very similar to the one in gauge invariant formulation of massive fields. Stueckelberg fields provide interesting possibilities for the study of interacting massive gauge fields (see e.g. Refs.[27, 28]). So we think that application of our approach to the interacting conformal self-dual fields may lead to new interesting development.

iii) BRST approach is one of powerful approaches to the analysis of various aspects of relativistic dynamics (see e.g. Refs.[29, 30]). This approach is conveniently adapted for the ordinary-derivative formulation. Recent application of BRST approach to the study of totally antisymmetric fields may be found in [31]. We think that extension of this approach to the case of conformal self-dual fields should be relatively straightforward.

iv) Self-dual fields studied in this paper are the particular case of mixed-symmetry fields. In the last years, there were interesting developments in the studying mixed-symmetry fields [32]-[35] that are invariant with respect to Poincaré algebra symmetries. It would be interesting to apply methods developed in Refs.[32]-[35] to the studying conformal self-dual mixed-symmetry fields⁸.

⁸ Unfolded form of equations of motion for conformal mixed-symmetry fields is studied in Ref.[36]. Higher-derivative Lagrangian formulation of the mixed-symmetry conformal fields was recently developed in Ref.[37].

There are other various interesting approaches in the literature which could be used to discuss the ordinary-derivative formulation of conformal self-dual fields. This is to say that various recently developed interesting formulations in terms of unconstrained fields in flat space may be found in Refs.[38, 39].

Acknowledgments. This work was supported by the RFBR Grant No.08-02-00963, RFBR Grant for Leading Scientific Schools, Grant No. 1615.2008.2, by the Dynasty Foundation and by the Alexander von Humboldt Foundation Grant PHYS0167.

Appendix A Derivation of ordinary-derivative gauge invariant Lagrangian

Because the methods for finding the ordinary-derivative Lagrangian for arbitrary $d \geq 4$ are quite similar we present details of the derivation for the case of $d = 4$. To derive ordinary-derivative gauge invariant Lagrangian (3.5) we use Lagrangian of the standard formulation given in (2.33). First, in place of the field T^{ab} , we introduce the fields t^{ab} and ϕ_{-1}^a by the relation

$$T^{ab} = t^{ab} + F^{ab}(\phi_{-1}) + \frac{\tau}{2}\epsilon^{abce}F^{ce}(\phi_{-1}), \quad (\text{A.1})$$

where t^{ab} satisfies self-duality constraint (3.2). Plugging (A.1) in (2.33) we obtain

$$\begin{aligned} \mathcal{L}_{\text{st}} = & -\frac{1}{2}F^{ab}(\bar{\phi}_{-1})\square F^{ab}(\phi_{-1}) \\ & - \frac{1}{2}\bar{t}^{ab}\square F^{ab}(\phi_{-1}) - \frac{1}{2}F^{ab}(\bar{\phi}_{-1})\square t^{ab} + \partial^a\bar{t}^{ac}\partial^bt^{bc}, \end{aligned} \quad (\text{A.2})$$

where the field strength $F^{ab}(\phi_{-1})$ is defined as in (3.6). We note that representation for T^{ab} given in (A.1) implies that T^{ab} is invariant under the gauge transformations

$$\delta\phi_{-1}^a = \partial^a\xi_{-2} - \lambda^a, \quad (\text{A.3})$$

$$\delta t^{ab} = \partial^a\lambda^b - \partial^b\lambda^a + \tau\epsilon^{abce}\partial^c\lambda^e. \quad (\text{A.4})$$

This implies that Lagrangian (A.2) is also invariant under gauge transformations (A.3), (A.4).

Second, we introduce new fields ϕ_1^a and ϕ_0 by using the following Lagrangian in place of (A.2)

$$\mathcal{L} = \mathcal{L}_{\text{st}} - \bar{X}^a X^a, \quad (\text{A.5})$$

where we use the notation

$$X^a \equiv \phi_1^a + \partial^a\phi_0 - \partial^b F^{ba}(\phi_{-1}) - \partial^b t^{ba}, \quad (\text{A.6})$$

and $\bar{X}^a$ is complex conjugate of X^a . It is clear that, on-shell, Lagrangians $\mathcal{L}_{\text{st}}$ and $\mathcal{L}$ describe the same field D.o.F. Using the formula (up to total derivative)

$$\partial^a t^{ac}\partial^b F^{bc} = -\frac{1}{2}t^{ab}\square F^{ab}, \quad (\text{A.7})$$

it is easy to see that Lagrangian (A.5) gives ordinary-derivative Lagrangian (3.5).

We now consider gauge symmetries of Lagrangian (A.5). Because the contribution to X^a given by (see (A.6))

$$- \partial^b F^{ba}(\phi_{-1}) - \partial^b t^{ba} \quad (\text{A.8})$$

is invariant under gauge transformations (A.3),(A.4), the X^a is also invariant under these gauge transformations, i.e., Lagrangian (A.5) is invariant under gauge transformations (A.3),(A.4). Besides this, we note that X^a is invariant under the additional gauge transformations

$$\delta\phi_1^a = \partial^a \xi_0, \quad \delta\phi_0 = -\xi_0. \quad (\text{A.9})$$

Altogether, gauge transformations (A.3),(A.4),(A.9) amount to the ones given in (3.10)-(3.13).

Appendix B Derivation of operator R^a

In this Appendix, we outline the derivation of the operator R^a (5.41). Use of the oscillator formulation turns out to be convenient for this purpose. The operator R^a is then determined by requiring action of self-dual field (4.4) to be invariant under the conformal boost transformations. In order to analyze restrictions imposed on the operator R^a by the conformal boost symmetries we need to know explicit form of the restrictions imposed on operator E (5.13) by the Lorentz and dilatation symmetries. Requiring action (4.4) with Lagrangian (5.12) to be invariant under the Lorentz and dilatation symmetries $\delta_{J^{ab}}S = 0$, $\delta_D S = 0$, amounts to the following respective equations for the operator E :

$$[E, J^{ab}] = 0, \quad [E, D] = 2E, \quad (\text{B.1})$$

where the operators J^{ab} and D are given in (2.17) and (2.18) respectively.

Variation of Lagrangian (5.12) under the conformal boost transformation can be presented as (up to total derivative)

$$\delta_{K^a} \mathcal{L} = \frac{\hbar}{2} \langle \Phi | \delta_{K^a} E | \Phi \rangle, \quad (\text{B.2})$$

$$\delta_{K^a} E \equiv K^{a\dagger} E + E K^a. \quad (\text{B.3})$$

Using (B.1) and K^a given in (2.19), we make sure that $\delta_{K^a} E$ (B.3) can be represented as

$$\delta_{K^a} E = R^{a\dagger} E + E R^a + E^a, \quad (\text{B.4})$$

$$E^a \equiv (\Delta + 1)[E, x^a] + [E, x^b] M^{ab} - \frac{1}{2} [[E, x^b], x^b] \partial^a, \quad (\text{B.5})$$

where M^{ab} and Δ are given in (5.38),(5.39). From (B.2), (B.4), we see that the requirement of invariance of the action under the conformal boost transformations amounts to the equations

$$R^{a\dagger} E + E R^a + E^a \approx 0. \quad (\text{B.6})$$

In (B.6) and below, to simplify our formulas, we adopt the following convention. Let A be some operator. We use the relation $A \approx 0$ in place of $\langle \Phi | A | \Phi \rangle = 0$.

Using operator E (5.13)-(5.19) and formula (B.5), we find immediately the operator E^a ,

$$E^a = \begin{pmatrix} E_{\phi\phi(1)}^a + E_{\phi\phi(0)}^a & -v^\ominus \bar{\alpha}^a \\ -\alpha^a \bar{v}^\ominus & 0 \end{pmatrix}, \quad (\text{B.7})$$

$$E_{\phi\phi(1)}^a = 2\Delta'\partial^a - (\Delta' + N_\zeta)\alpha^a\bar{\alpha}\partial + (-\Delta' + N_\zeta)\alpha\partial\bar{\alpha}^a, \quad (\text{B.8})$$

$$E_{\phi\phi(0)}^a = (\Delta' - N_\zeta)e_1\bar{\alpha}^a + \alpha^a\bar{e}_1(\Delta' + N_\zeta), \quad (\text{B.9})$$

where e_1 , $\bar{e}_1$ and Δ' are given in (5.19) and (5.39) respectively. Also, we note that the commutation relation $[D, K^a] = K^a$ gives the following equation for the operator R^a :

$$[D, R^a] = R^a. \quad (\text{B.10})$$

Equations (B.6), (B.10) constitute a complete system of equations which allows us to determine the operator R^a uniquely. We now discuss the procedure of solving these equations.

Operator E (5.13) is a second-order polynomial in the derivative. From (B.7), we see that the operator E^a is a first-order polynomial in the derivative. The operator R^a is also turned out to be first-order polynomial in the derivative. Therefore it is convenient to represent the operators E , E^a , and R^a as power series in the derivative,

$$E = E_{(2)} + E_{(1)} + E_{(0)}, \quad (\text{B.11})$$

$$E^a = E_{(1)}^a + E_{(0)}^a, \quad R^a = R_{(1)}^a + R_{(0)}^a, \quad (\text{B.12})$$

where the operators $E_{(n)}$, $E_{(n)}^a$, and $R_{(n)}^a$ are degree- n homogeneous polynomials in the derivative. Explicit expressions for the operators $E_{(n)}$ and $E_{(n)}^a$ can easily be read from the respective expressions in (5.13)-(5.19) and (B.7)-(B.9). Using (B.11), (B.12) it is easy to see that Eqs.(B.6) amount to the following equations:

$$E_{(2)}R_{(1)}^a + h.c. \approx 0, \quad (\text{B.13})$$

$$E_{(2)}R_{(0)}^a + E_{(1)}R_{(1)}^a + h.c. \approx 0, \quad (\text{B.14})$$

$$\left(E_{(1)}R_{(0)}^a + E_{(0)}R_{(1)}^a + h.c.\right) + E_{(1)}^a \approx 0, \quad (\text{B.15})$$

$$\left(E_{(0)}R_{(0)}^a + h.c.\right) + E_{(0)}^a \approx 0. \quad (\text{B.16})$$

We now present our procedure for solving Eqs.(B.10) and (B.13)-(B.16).

i) We note that the most general operators $R_{(n)}^a$ (B.12) acting on 2-vector $|\Phi\rangle$ (5.1) can be presented as 2×2 matrices given by

$$R_{(n)}^a = \begin{pmatrix} R_{\phi\phi(n)}^a & R_{\phi t(n)}^a \\ R_{t\phi(n)}^a & R_{tt(n)}^a \end{pmatrix}, \quad n = 0, 1. \quad (\text{B.17})$$

Requiring the operator R^a to satisfy Eq.(B.10) and constraints (5.28)-(5.31) we find the following

expressions⁹

$$R_{\phi\phi(0)}^a = r_{0,1}\bar{\alpha}^a + \alpha^a\bar{r}_{0,1}, \quad (\text{B.18})$$

$$R_{\phi\phi(1)}^a = r_{1,1}\partial^a + r_{1,5}\alpha^a\bar{\alpha}\partial + r_{\epsilon 1}\epsilon_0^{ab}\partial^b, \quad (\text{B.19})$$

$$R_{\phi t(0)}^a = r_{0,4}\bar{\alpha}^a, \quad (\text{B.20})$$

$$R_{\phi t(1)}^a = 0, \quad (\text{B.21})$$

$$R_{t\phi(0)}^a = r_{0,5}(1 + \epsilon)\alpha^a, \quad (\text{B.22})$$

$$R_{t\phi(1)}^a = 0, \quad (\text{B.23})$$

$$R_{tt(n)}^a = 0, \quad n = 0, 1, \quad (\text{B.24})$$

where the operators $r_{0,1}, r_{0,2}, r_{0,4}, r_{0,5}, r_{1,1}, r_{1,5}, r_{\epsilon 1}$ independent of the oscillators α^a are given by

$$r_{0,1} = \zeta\tilde{r}_{0,1}\bar{v}^\oplus, \quad \bar{r}_{0,1} = v^\oplus\tilde{\bar{r}}_{0,1}\bar{\zeta}, \quad (\text{B.25})$$

$$r_{1,1} = v^\oplus\tilde{r}_{1,1}\bar{v}^\oplus, \quad r_{\epsilon,1} = v^\oplus\tilde{r}_{\epsilon,1}\bar{v}^\oplus, \quad r_{1,5} = v^\oplus\tilde{r}_{1,5}\bar{v}^\oplus, \quad (\text{B.26})$$

$$r_{0,4} = v^\oplus\tilde{r}_{0,4}, \quad r_{0,5} = \tilde{r}_{0,5}\bar{v}^\oplus, \quad (\text{B.27})$$

and the quantities $\tilde{r}_{0,1}, \tilde{\bar{r}}_{0,1}, \tilde{r}_{0,4}, \tilde{r}_{0,5}, \tilde{r}_{1,1}, \tilde{r}_{1,5}, \tilde{r}_{\epsilon 1}$ independent of the oscillators $\alpha^a, \zeta, v^\oplus, v^\ominus$ remain as undermined constants. Below, we determine these quantities by using Eqs.(B.13)-(B.16).

Before analyzing Eqs.(B.13)-(B.16) we explain our terminology. Introducing the notation $\mathcal{X}$ for the left hand side of Eqs.(B.13)-(B.16) we note that $\mathcal{X}$ is a 2×2 matrix acting on 2-vector $|\Phi\rangle$ (5.1). Using the notation

$$\mathcal{X} = \begin{pmatrix} \mathcal{X}_{\phi\phi} & \mathcal{X}_{\phi t} \\ \mathcal{X}_{t\phi} & \mathcal{X}_{tt} \end{pmatrix}, \quad (\text{B.28})$$

we note that the 2×2 matrix equation $\mathcal{X} \approx 0$ amounts to the four equations, $\langle\phi|\mathcal{X}_{\phi\phi}|\phi\rangle = 0$, $\langle\phi|\mathcal{X}_{\phi t}|t\rangle = 0$, $\langle t|\mathcal{X}_{t\phi}|\phi\rangle = 0$, $\langle t|\mathcal{X}_{tt}|t\rangle = 0$. We refer to these four equations to as the respective $\phi\phi$ -, ϕt -, $t\phi$ -, and tt -parts of the equation $\mathcal{X} \approx 0$. We now turn to analysis of Eqs.(B.13)-(B.16).

ii) Using $\phi\phi$ part of Eq.(B.13) and the relation

$$E_{\phi\phi(2)}R_{\phi\phi(1)}^a = r_{1,1}E_{\phi\phi(2)}\partial^a + r_{1,5}\square\alpha^a\bar{\alpha}\partial - r_{1,5}\alpha\partial\bar{\alpha}\partial\partial^a + r_{\epsilon,1}\square\epsilon_0^{ab}\partial^b, \quad (\text{B.29})$$

we find

$$\tilde{r}_{1,1}^\dagger = \tilde{r}_{1,1}, \quad \tilde{r}_{\epsilon,1}^\dagger = \tilde{r}_{\epsilon,1}, \quad \tilde{r}_{1,5} = 0. \quad (\text{B.30})$$

iii) Making use of $\phi\phi$ part of Eq.(B.14) and the relations

$$E_{\phi\phi(2)}R_{\phi\phi(0)}^a = \alpha^a E_{\phi\phi(2)}\bar{r}_{0,1} + r_{0,1}E_{\phi\phi(2)}\bar{\alpha}^a - \alpha\partial\partial^a\bar{r}_{0,1}, \quad (\text{B.31})$$

$$E_{\phi\phi(1)}R_{\phi\phi(1)}^a = e_1 r_{1,1}\bar{\alpha}\partial\partial^a + \alpha\partial\partial^a\bar{e}_1 r_{1,1}, \quad (\text{B.32})$$

we obtain

$$\bar{r}_{0,1} = [\bar{e}_1, r_{1,1}], \quad \tilde{r}_{0,1}^\dagger = -\tilde{\bar{r}}_{0,1}. \quad (\text{B.33})$$

⁹ As a realization of the operator R^a on the gauge field $|\Phi\rangle$ is defined up to gauge transformation (5.32) we ignore contributions to R^a that can be removed by gauge transformation (5.32).

iv) Using $\phi\phi$ part of Eq.(B.15) and the relations

$$E_{\phi\phi(1)}R_{\phi\phi(0)}^a = e_1\bar{r}_{0,1}\partial^a - e_1\bar{r}_{0,1}\alpha^a\bar{\alpha}\partial + \bar{e}_1r_{0,1}\alpha\partial\bar{\alpha}^a, \quad (\text{B.34})$$

$$E_{\phi\phi(0)}R_{\phi\phi(1)}^a = m_1r_{1,1}\partial^a + m_1r_{\epsilon,1}\epsilon_0^{ab}\partial^b, \quad (\text{B.35})$$

$$E_{\phi t}R_{t\phi(0)}^a = v^\ominus r_{0,5}\partial^a - v^\ominus r_{0,5}\alpha^a\bar{\alpha}\partial - v^\ominus r_{0,5}\epsilon_0^{ab}\partial^b, \quad (\text{B.36})$$

we find

$$\tilde{r}_{0,5} = 1, \quad \tilde{r}_{\epsilon,1} = -1, \quad \tilde{r}_{0,1} = 1, \quad \tilde{r}_{1,1} = -1. \quad (\text{B.37})$$

v) Using ϕt part of Eq.(B.14) and the relation

$$E_{\phi\phi(2)}R_{\phi t(0)}^a + (E_{t\phi}R_{\phi\phi(1)}^a)^\dagger = (r_{0,4} - r_{1,1}v^\ominus)\bar{\alpha}\partial + (r_{0,4} - r_{\epsilon,1}v^\ominus)\epsilon_0^{ab}\partial^b\bar{\alpha}\partial \quad (\text{B.38})$$

we obtain

$$\tilde{r}_{0,4} = -1, \quad \tilde{r}_{0,4} = \tilde{r}_{\epsilon,1}. \quad (\text{B.39})$$

vi) Using $\tilde{r}_{0,1}, \tilde{r}_{0,1}, \tilde{r}_{0,4}, \tilde{r}_{0,5}, \tilde{r}_{1,1}, \tilde{r}_{1,5}, \tilde{r}_{\epsilon,1}$ above-given, we make sure that all the remaining equations in (B.13)-(B.16) are satisfied automatically. We note that in analysis of tt part of Eq.(B.15) we use the identity

$$\langle t|\alpha^a\bar{\alpha}\partial|t\rangle = \langle t|\alpha\partial\bar{\alpha}^a|t\rangle, \quad (\text{B.40})$$

which can be proved by using self-duality constraint (5.10).

Appendix C On-shell D.o.F of self-dual field in $4d$

We analyze on-shell D.o.F of the conformal self-dual field in $4d$ with Lagrangian (3.5). To this end we use light-cone gauge. In light-cone frame, the space-time coordinates x^a are decomposed as $x^a = x^+, x^-, x^i$, where the light-cone coordinates in $\pm$ directions are defined as $x^\pm = (x^3 \pm x^0)/\sqrt{2}$ and x^+ is taken to be a light-cone time. The $so(2)$ algebra vector indices take values $i, j = 1, 2$. We adopt the conventions: $\partial^i = \partial_i \equiv \partial/\partial x^i$, $\partial^\pm = \partial_\mp \equiv \partial/\partial x^\mp$.

We are going to prove that on-shell D.o.F of the self-dual field are described by two $so(2)$ algebra self-dual complex-valued vector fields ϕ_{-1}^i, ϕ_1^i and one complex-valued scalar field ϕ_0 ,

$$\phi_{-1}^i, \quad \phi_1^i, \quad \phi_0, \quad (\text{C.1})$$

which satisfy the equations of motion

$$\square\phi_{-1}^i - \phi_1^i = 0, \quad \square\phi_1^i = 0, \quad \square\phi_0 = 0. \quad (\text{C.2})$$

The vector fields satisfy the $so(2)$ self-duality constraint:

$$\pi^{ij}\phi_{-1}^j = 0, \quad \pi^{ij}\phi_1^j = 0. \quad (\text{C.3})$$

Here and below, we use the notation

$$\pi^{ij} \equiv \delta^{ij} + \tau\epsilon^{ij}, \quad \bar{\pi}^{ij} \equiv \delta^{ij} - \tau\epsilon^{ij}, \quad (\text{C.4})$$

where δ^{ij} is the Kronecker delta, while ϵ^{ij} is the Levi-Civita symbol normalized as $\epsilon^{12} = 1$.

In order to find on-shell D.o.F we use equations of motion obtained from Lagrangian (3.5) and self-duality constraint (3.2),

$$\partial^a F^{ab}(\phi_1) = 0, \quad (C.5)$$

$$\partial^b F^{ba}(\phi_{-1}) + \partial^b t^{ba} - \phi_1^a - \partial^a \phi_0 = 0, \quad (C.6)$$

$$\square \phi_0 + \partial^a \phi_1^a = 0, \quad (C.7)$$

$$F^{ab}(\phi_1) = \frac{\tau}{2} \epsilon^{abce} F^{ce}(\phi_1), \quad (C.8)$$

$$t^{ab} = \frac{\tau}{2} \epsilon^{abce} t^{ce}. \quad (C.9)$$

Taking into account the light-cone frame decomposition of the vector and tensor fields (3.1),

$$\phi_{-1}^a = \phi_{-1}^+, \phi_{-1}^-, \phi_{-1}^i, \quad \phi_1^a = \phi_1^+, \phi_1^-, \phi_1^i, \quad t^{ab} = t^{+-}, t^{+i}, t^{-i}, t^{ij}, \quad (C.10)$$

we note that some of gauge transformations given (3.10)-(3.13) can be represented as

$$\delta \phi_1^+ = \partial^+ \xi_0, \quad (C.11)$$

$$\delta \phi_{-1}^+ = \partial^+ \xi_{-2} - \lambda^+, \quad (C.12)$$

$$\delta \phi_{-1}^i = \partial^i \xi_{-2} - \lambda^i, \quad (C.13)$$

$$\delta t^{+-} = \partial^+ \lambda^- - \partial^- \lambda^+ + \tau \epsilon^{ij} \partial^i \lambda^j, \quad (C.14)$$

$$\delta t^{+i} = \bar{\pi}^{ij} (\partial^+ \lambda^j - \partial^j \lambda^+). \quad (C.15)$$

From (C.11), we see that the field ϕ_1^+ can be gauge away by using ξ_0 gauge transformation. From (C.12), (C.13), we see that the fields ϕ_{-1}^+ and $\pi^{ij} \phi_{-1}^j$ can be gauged away by using the respective ξ_{-2} and $\pi^{ij} \lambda^j$ gauge transformations. From (C.14), (C.15), we see that the fields t^{+-} and $\bar{\pi}^{ij} t^{+j}$ can be gauged away by using the respective λ^- and $\bar{\pi}^{ij} \lambda^j$ gauge transformations. To summarize, we can impose the following gauge conditions:

$$\phi_1^+ = 0, \quad (C.16)$$

$$\phi_{-1}^+ = 0, \quad (C.17)$$

$$\pi^{ij} \phi_{-1}^j = 0, \quad (C.18)$$

$$\bar{\pi}^{ij} t^{+j} = 0, \quad (C.19)$$

$$t^{+-} = 0. \quad (C.20)$$

Using gauge conditions (C.16)-(C.20), one can make sure that equations (C.5)-(C.9) amount to the following equations

$$\square \phi_1^a = 0, \quad (C.21)$$

$$\partial^a \phi_1^a = 0, \quad (C.22)$$

$$\partial^a \phi_{-1}^a + \phi_0 = 0. \quad (C.23)$$

$$\square \phi_{-1}^a + \partial^b t^{ba} - \phi_1^a = 0, \quad (C.24)$$

$$\square \phi_0 = 0, \quad (C.25)$$

$$\pi^{ij}\phi_1^j = 0, \quad (\text{C.26})$$

$$\pi^{ij}t^{+j} = 0, \quad (\text{C.27})$$

$$\bar{\pi}^{ij}t^{-j} = 0, \quad (\text{C.28})$$

$$t^{ij} = 0. \quad (\text{C.29})$$

We now analyze gauge conditions (C.16)-(C.20) and equations (C.21)-(C.29).

i) In view of (C.18), (C.26), we see that the $so(2)$ algebra vector fields ϕ_{-1}^i , ϕ_1^i are indeed satisfy the self-duality constraint given in (C.3).

ii) Equation (C.21) leads to the 2nd equation in (C.2).

iii) Differential constraints (C.22), (C.23) and gauge conditions (C.16), (C.17) tell us that the non-dynamical fields ϕ_1^- and ϕ_{-1}^- can be expressed in terms of fields (C.1),

$$\phi_1^- = -\frac{\partial^i}{\partial^+}\phi_1^i, \quad \phi_{-1}^- = -\frac{\partial^i}{\partial^+}\phi_{-1}^i - \frac{1}{\partial^+}\phi_0. \quad (\text{C.30})$$

iii) Equations (C.19), (C.27) imply

$$t^{+i} = 0. \quad (\text{C.31})$$

iv) Taking into account (C.29), (C.31) and using Eq.(C.24) we obtain

$$\square\phi_{-1}^i + \partial^+t^{-i} - \phi_1^i = 0. \quad (\text{C.32})$$

Multiplying Eq.(C.32) by $\bar{\pi}^{ij}$ and using constraint (C.18) we obtain the 1st equation in (C.2).

v) Multiplying Eq.(C.32) by π^{ij} and taking into account (C.18),(C.26) gives the equation

$$\pi^{ij}t^{-j} = 0. \quad (\text{C.33})$$

Equations (C.28),(C.33) imply

$$t^{-i} = 0. \quad (\text{C.34})$$

Taking into account (C.20), (C.29), (C.31), (C.34), we see that $t^{ab} = 0$.

To summarize, we proved that on-shell D.o.F of the self-dual field with Lagrangian (3.5) are described by fields given in (C.1). These fields satisfy equations of motion (C.2) and self-duality constraints (C.3). Light-cone gauge Lagrangian which leads to equations of motion (C.2) is given in (3.39).

Appendix D ϵ -symbols

In this Appendix, we describe various useful relations for ϵ -symbols we use in the paper. We introduce the following ϵ -symbols constructed out of the Levi-Civita symbol $\epsilon^{a_1\dots a_\nu b_1\dots b_\nu}$ and the oscillators:

$$\epsilon \equiv \frac{\tau}{(\nu!)^2} \epsilon^{a_1 \dots a_\nu b_1 \dots b_\nu} \alpha^{a_1} \dots \alpha^{a_\nu} \bar{\alpha}^{b_\nu} \dots \bar{\alpha}^{b_1}, \quad (\text{D.1})$$

$$\epsilon^a \equiv \frac{\tau}{\nu!(\nu-1)!} \epsilon^{a_1 \dots a_\nu a b_2 \dots b_\nu} \alpha^{a_1} \dots \alpha^{a_\nu} \bar{\alpha}^{b_\nu} \dots \bar{\alpha}^{b_2}, \quad (\text{D.2})$$

$$\bar{\epsilon}^a \equiv \frac{\tau}{\nu!(\nu-1)!} \epsilon^{a a_2 \dots a_\nu b_1 \dots b_\nu} \alpha^{a_2} \dots \alpha^{a_\nu} \bar{\alpha}^{b_\nu} \dots \bar{\alpha}^{b_1}, \quad (\text{D.3})$$

$$\epsilon_0^{ab} \equiv \frac{\tau}{((\nu-1)!)^2} \epsilon^{a a_2 \dots a_\nu b b_2 \dots b_\nu} \alpha^{a_2} \dots \alpha^{a_\nu} \bar{\alpha}^{b_\nu} \dots \bar{\alpha}^{b_2}, \quad (\text{D.4})$$

$$\epsilon^{ab} \equiv \frac{\tau}{\nu!(\nu-2)!} \epsilon^{a_1 \dots a_\nu a b b_3 \dots b_\nu} \alpha^{a_1} \dots \alpha^{a_\nu} \bar{\alpha}^{b_\nu} \dots \bar{\alpha}^{b_3}, \quad (\text{D.5})$$

$$\bar{\epsilon}^{ab} \equiv \frac{\tau}{\nu!(\nu-2)!} \epsilon^{a b a_3 \dots a_\nu b_1 \dots b_\nu} \alpha^{a_3} \dots \alpha^{a_\nu} \bar{\alpha}^{b_\nu} \dots \bar{\alpha}^{b_1}, \quad (\text{D.6})$$

where τ is defined in (2.28). We note the following helpful relations for these ϵ -symbols:

$$\epsilon^a \equiv [\epsilon, \alpha^a], \quad \bar{\epsilon}^a \equiv [\bar{\alpha}^a, \epsilon], \quad (\text{D.7})$$

$$\epsilon^{ab} \equiv \{[\epsilon, \alpha^a], \alpha^b\}, \quad \bar{\epsilon}^{ab} \equiv \{\bar{\alpha}^a, [\bar{\alpha}^b, \epsilon]\}, \quad (\text{D.8})$$

$$\begin{aligned} \epsilon_0^{ab} &\equiv \{[\bar{\alpha}^a, \epsilon], \alpha^b\}, \\ &= \{\bar{\alpha}^a, \epsilon^b\} = \{\bar{\epsilon}^a, \alpha^b\}. \end{aligned} \quad (\text{D.9})$$

Our ϵ -symbols satisfy the following hermitian conjugation rules:

$$\epsilon^\dagger = -\epsilon, \quad \epsilon^{a\dagger} = -\bar{\epsilon}^a, \quad \epsilon_0^{ab\dagger} = \epsilon_0^{ab}, \quad \epsilon^{ab\dagger} = \bar{\epsilon}^{ab}. \quad (\text{D.10})$$

On space of ket-vector $|\phi\rangle$ subject to the constraint

$$N_\alpha |\phi\rangle = \nu |\phi\rangle, \quad (\text{D.11})$$

we obtain the relation

$$\epsilon^2 |\phi\rangle = |\phi\rangle. \quad (\text{D.12})$$

It is this property of the ϵ -symbol that is used for the definition of the self-dual ket-vector $|t\rangle$ (5.10).

One has the following helpful identities involving ϵ -symbols, the oscillators, and the derivative

$$\alpha \partial \epsilon_0^{ab} \partial^b = \epsilon^b \partial^b \partial^a - \epsilon^a \square + \epsilon^{ab} \bar{\alpha} \partial \partial^b, \quad (\text{D.13})$$

$$\epsilon_0^{ab} \bar{\alpha} \partial \partial^b = -\bar{\epsilon}^b \partial^b \partial^a + \bar{\epsilon}^a \square + \alpha \partial \bar{\epsilon}^{ab} \partial^b. \quad (\text{D.14})$$

References

- [1] J. H. Schwarz and P. C. West, Phys. Lett. B **126**, 301 (1983).
- [2] N. Marcus and J. H. Schwarz, Phys. Lett. B **115**, 111 (1982).
- [3] N. Berkovits, Phys. Lett. B **395**, 28 (1997) [arXiv:hep-th/9610134].

- [4] P. Pasti, D. P. Sorokin and M. Tonin, Phys. Rev. D **55**, 6292 (1997) [arXiv:hep-th/9611100].
- [5] I. Bandos, X. Bekaert, J. A. de Azcarraga, D. Sorokin and M. Tsulaia, JHEP **0505**, 031 (2005) [arXiv:hep-th/0501113].
- [6] E. S. Fradkin and A. A. Tseytlin, Phys. Rept. **119**, 233 (1985).
- [7] Yu. M. Zinoviev, “On massive high spin particles in (A)dS,” arXiv:hep-th/0108192.
- [8] R. R. Metsaev, Phys. Lett. B **643**, 205 (2006) [arXiv:hep-th/0609029].
- [9] V. E. Lopatin and M. A. Vasiliev, Mod. Phys. Lett. A **3**, 257 (1988).
- [10] M. A. Vasiliev, Nucl. Phys. B **301**, 26 (1988).
- [11] J. M. F. Labastida, Nucl. Phys. B **322**, 185 (1989).
- [12] R. R. Metsaev, Phys. Lett. B **354**, 78 (1995).
- [13] X. Bekaert and N. Boulanger, Commun. Math. Phys. **245**, 27 (2004) [arXiv:hep-th/0208058].
- [14] X. Bekaert and N. Boulanger, Commun. Math. Phys. **271**, 723 (2007) [arXiv:hep-th/0606198].
- [15] K. Hallowell and A. Waldron, Nucl. Phys. B **724**, 453 (2005) [arXiv:hep-th/0505255].
- [16] A. Y. Artsukevich and M. A. Vasiliev, Phys. Rev. D **79**, 045007 (2009) [arXiv:0810.2065 [hep-th]].
- [17] J. Engquist and P. Sundell, Nucl. Phys. B **752**, 206 (2006) [arXiv:hep-th/0508124].
- [18] F. Bastianelli, O. Corradini and E. Latini, JHEP **0702**, 072 (2007) [arXiv:hep-th/0701055].
- [19] F. Bastianelli, O. Corradini and E. Latini, JHEP **0811**, 054 (2008) [arXiv:0810.0188 [hep-th]].
- [20] E. Bergshoeff, M. de Roo and B. de Wit, Nucl. Phys. B **182**, 173 (1981).
- [21] E. Bergshoeff, M. de Roo and B. de Wit, Nucl. Phys. B **217**, 489 (1983).
- [22] E. Bergshoeff, E. Sezgin and A. Van Proeyen, Nucl. Phys. B **264**, 653 (1986) [Erratum-ibid. B **598**, 667 (2001)].
- [23] E. Bergshoeff, E. Sezgin and A. Van Proeyen, Class. Quant. Grav. **16**, 3193 (1999) hep-th/9904085
- [24] E. Bergshoeff, A. Salam and E. Sezgin, Nucl. Phys. B **279**, 659 (1987).
- [25] R. R. Metsaev, “Ordinary-derivative formulation of conformal low spin fields,” arXiv:0707.4437 [hep-th]. “Ordinary-derivative formulation of conformal totally symmetric arbitrary spin bosonic fields,” arXiv:0709.4392 [hep-th].
- [26] R. R. Metsaev, Phys. Rev. D **78**, 106010 (2008) [arXiv:0805.3472 [hep-th]].
- [27] Yu. M. Zinoviev, Nucl. Phys. B **770**, 83 (2007) [arXiv:hep-th/0609170].
- [28] R. R. Metsaev, Phys. Rev. D **77**, 025032 (2008) [arXiv:hep-th/0612279].
- [29] W. Siegel, “Fields,” arXiv:hep-th/9912205.
- [30] A. Fotopoulos and M. Tsulaia, Int. J. Mod. Phys. A **24**, 1 (2009) [arXiv:0805.1346 [hep-th]].
- [31] I. L. Buchbinder, V. A. Krykhtin and L. L. Ryskina, Mod. Phys. Lett. A **24**, 401 (2009) [arXiv:0810.3467 [hep-th]].
- [32] Yu. M. Zinoviev, “On massive mixed symmetry tensor fields in Minkowski space and (A)dS,” arXiv:hep-th/0211233. “First order formalism for mixed symmetry tensor fields,” arXiv:hep-th/0304067. “First order formalism for massive mixed symmetry tensor fields in arXiv:hep-th/0306292.
- [33] E. D. Skvortsov, JHEP **0807**, 004 (2008) [arXiv:0801.2268 [hep-th]]. Nucl. Phys. B **808**, 569 (2009) [arXiv:0807.0903 [hep-th]].
- [34] K. B. Alkalaev, M. Grigoriev and I. Y. Tipunin, Nucl. Phys. B **823**, 509 (2009) [arXiv:0811.3999 [hep-th]].
- [35] P. Y. Moshin and A. A. Reshetnyak, JHEP **0710**, 040 (2007) [arXiv:0707.0386 [hep-th]].
- [36] O. V. Shaynkman, I. Y. Tipunin and M. A. Vasiliev, Rev. Math. Phys. **18**, 823 (2006) hep-th/0401086
- [37] M. A. Vasiliev, “Bosonic conformal higher-spin fields of any symmetry,” arXiv:0909.5226 [hep-th].
- [38] I. L. Buchbinder and A. V. Galajinsky, JHEP **0811**, 081 (2008) [arXiv:0810.2852 [hep-th]].
- [39] A. Campoleoni, D. Francia, J. Mourad and A. Sagnotti, Nucl. Phys. B **815**, 289 (2009) [arXiv:0810.4350 [hep-th]].